

April Forecast Update for Northwest Pacific Typhoon Activity in 2003

Issued: 11th April 2003

by Drs Adam Lea and Mark Saunders Benfield Hazard Research Centre, UCL (University College London), UK

Forecast Summary

TSR anticipates the 2003 Northwest Pacific typhoon season will see activity slightly above average.

The TSR (Tropical Storm Risk) early April forecast update for Northwest Pacific typhoon activity in 2003 anticipates a slightly above average season. The forecast spans the full Northwest Pacific season from 1st January to 31st December 2003 and is based on data available through the end of March 2003. Our predictor is the forecast anomaly in August-September Niño 4 sea surface temperature (SST) which we anticipate will be $0.20\pm0.39^{\circ}$ C warmer than normal this summer. Monthly updated forecasts will be issued through to early August. This month we introduce the percentage improvement in mean square error over a climatological forecast as our skill score, this being the standard skill metric recommended by the WMO.

NW Pacific ACE Index and System Numbers in 2003

				ACE Index	Intense Typhoons	Typhoons	Tropical Storms		
TSR Forecast (±FE)		2003	318 (±102)	9.2 (±2.9)	17.1 (±4.5)	26.7 (±5.1)			
10yr Climate Norm (±SD)		1993-2002	300 (±113)	9.1 (±3.2)	17.2 (±4.7)	27.8 (±5.0)			
30yr Climate Norm (±SD)			1973-2002	285 (±97)	8.0 (±3.0)	16.6 (±3.7)	26.3 (±4.3)		
Forecast Skill at this Lead		1988-2002	25%	17%	15%	3%			
Key:	ey: ACE Index = Accumulated Cyclone Energy Index = Sum of the Squares of 6-hourly Maximu Sustained Wind Speeds (in units of knots) for all Systems while they are at least Tropic Storm Strength. ACE Unit = x10 ⁴ knots ² .								
	Intense Typhoon	e Typhoon = 1 Minute Sustained Wind > 95Kts = Hurricane Category 3 to 5.							
	Typhoon = 1 Minute Sustained Wind > 63Kts = Hurricane Category 1 to 5.								
Tropical Storm = 1 Minute Sustained Wind > 33Kts.									
	SD	=	Standard Deviation.						
FE (Forecast Error) = Standard Deviation of Errors in Replicated Real Time Forecasts 1993-2002						002			
	Forecast Skill = Percentage Improvement in Mean Square Error over Running 10-year Prior Climat								
	Northwest Pacific	=	Northern Hemisph				a Sea Any		
Northwest Pacific = Northern Hemisphere Region West of 180°W Including the Tropical Cyclone (Irrespective of Where it Forms) Which I Strength Within this Region Counts as an Event.						-	-		

Key Predictor for 2003

The key factor behind our forecast for a slightly above average Northwest Pacific typhoon season in 2003 is the anticipated slightly positive Niño 4 (150°W-160°E, 5°S-5°N) SST anomaly. Our forecast anomaly (1973-2002 climatology) for August-September 2003 Niño 4 SST is 0.20±0.39°C (up from 0.01±0.42°C last month). The forecast skill for this predictor at this lead is 37%.

1

New Developments in 2003

For the 2003 Northwest Pacific typhoon season, TSR are: (1) Issuing monthly updated forecasts for basin tropical storm, typhoon and intense typhoon numbers and additionally for the ACE Index. The latter reflects a combination of intensity and duration for all storms each season and may be linked more closely to total losses and disruption than the individual number of tropical storms, typhoons or even intense typhoons; (2) Providing real-time forecasts through the TSR Storm Tracker for active Northwest Pacific storm systems. These forecasts are updated every 6-12 hours and provide the best available information on storm strength, track and track uncertainty all with various levels of zoom. A unique feature to be included soon is current and forecast 2-dimensional windfields for systems of at least typhoon force.

Further Information

Further information on the TSR forecast methodology, the TSR replicated real-time forecast skill 1987-2001 as a function of lead time, and on TSR in general, may be obtained from the 'Extended Range Forecast for Northwest Pacific Typhoon Activity in 2002' document issued on the 6th March 2002. Our next monthly forecast update for the 2003 Northwest Pacific typhoon season will be issued on the 6th May 2003. Further regular monthly updates will follow through to early August 2003.

Appendix - Predictions from Previous Months

NW Pacific ACE Index and System Numbers 2003									
		ACE Index	Tropical Storms	Typhoons	Intense Typhoons				
Average Number (±	SD) (1993-2002)	300 (±113)	27.8 (±5.0)	17.2 (±4.7)	9.1 (±3.2)				
Average Number (±	SD) (1973-2002)	285 (±97)	26.7 (±4.3)	16.6 (±3.7)	8.0 (±3.0)				
TSR Forecasts (±FE)	11th April 2003	318 (±102)	26.7 (±5.1)	17.1 (±4.5)	9.2 (±2.9)				
TSK Polecasts (TFE)	5th March 2003	297 (±100)	26.2 (±5.1)	16.6 (±4.5)	8.5 (±2.9)				

