

July Forecast Update for Northwest Pacific Typhoon Activity in 2003

Issued: 4th July 2003

by Drs Adam Lea and Mark Saunders Benfield Hazard Research Centre, UCL (University College London), UK

Forecast Summary

TSR slightly raises its forecast and anticipates the 2003 Northwest Pacific typhoon season will see activity close to average.

The TSR (Tropical Storm Risk) early July forecast update for Northwest Pacific typhoon activity in 2003 anticipates a near average season. The forecast spans the full Northwest Pacific season from 1st January to 31st December 2003 and is based on data available through the end of June 2003. The TSR predictor is the forecast anomaly in August-September Niño 4 sea surface temperature (SST) which we anticipate will be $0.03\pm0.24^{\circ}$ C warmer than normal this summer. The slight rise in forecast storm activity from last month comes from a warmer Niño 4 SST now being anticipated. The appendix gives the TSR predictions from previous months.

NW Pacific ACE Index and System Numbers in 2003

				ACE Index	Intense Typhoons	Typhoons	Tropical Storms		
TSR Forecast (±FE)			2003	299 (±85)	8.6 (±2.1)	16.7 (±4.1)	26.3 (±4.9)		
10yr Climate Norm (±SD)			1993-2002	300 (±113)	9.1 (±3.2)	17.2 (±4.7)	27.8 (±5.0)		
30yr Climate Norm (±SD)			1973-2002	285 (±97)	$8.0 (\pm 3.0)$	16.6 (±3.7)	26.3 (±4.3)		
Forecast Skill at this Lead			1988-2002	45%	57%	28%	14%		
·	Key: ACE Index = Accumulated Cyclone Energy Index = Sum of the Squares of 6-hourly Maximu Sustained Wind Speeds (in units of knots) for all Systems while they are at least Tropic Storm Strength. ACE Unit = x10 ⁴ knots ² . Intense Typhoon = 1 Minute Sustained Wind > 95Kts = Hurricane Category 3 to 5.								
	Typhoon	=	1 Minute Sustained Wind > 63Kts = Hurricane Category 1 to 5.						
Tropical Storm =		1 Minute Sustained Wind > 33Kts.							
	SD	=	Standard Deviation.						
]	FE (Forecast Error)	=	Standard Deviation of Errors in Replicated Real Time Forecasts 1993-2002						
Forecast Skill = Percentage Improvement in Mean Square Error over Running 10-year Prior C Norm from Replicated Real Time Forecasts 1988-2002. Northwest Pacific = Northern Hemisphere Region West of 180°W Including the South China Sea. Tropical Cyclone (Irrespective of Where it Forms) Which Reaches Tropical S Strength Within this Region Counts as an Event.							rior Climate		
						•			

Key Predictor for 2003

The key factor behind the TSR forecast for an average Northwest Pacific typhoon season in 2003 is the anticipated near neutral Niño 4 (150°W-160°E, 5°S-5°N) SST anomaly. The TSR forecast anomaly (1973-2002 climatology) for August-September 2003 Niño 4 SST is 0.03 ± 0.24 °C (up from -0.17±0.29°C last month). The forecast skill for this predictor at this lead is 81%.

1

Further Information

Further information on the TSR forecast methodology, the TSR replicated real-time forecast skill 1987-2001 as a function of lead time, and on TSR in general, may be obtained either from the 'Extended Range Forecast for Northwest Pacific Typhoon Activity in 2002' document issued on the 6th March 2002 or from the TSR web site tropicalstormrisk.com. The TSR next monthly forecast update for the 2003 Northwest Pacific typhoon season will be issued on the 5th August 2003.

Appendix - Predictions from Previous Months

NW Pacific ACE Index and System Numbers 2003									
		ACE Index	Tropical Storms	Typhoons	Intense Typhoons				
Average Number (±	SD) (1993-2002)	300 (±113)	27.8 (±5.0)	17.2 (±4.7)	9.1 (±3.2)				
Average Number (±	SD) (1973-2002)	285 (±97)	26.7 (±4.3)	16.6 (±3.7)	8.0 (±3.0)				
	4 Jul 2003	299 (±85)	26.3 (±4.9)	16.7 (±4.1)	8.6 (±2.1)				
	10 Jun 2003	275 (±89)	25.8 (±4.8)	16.1 (±4.1)	7.9 (±2.3)				
TSR Forecasts (±FE)	6 May 2003	284 (±84)	26.0 (±4.9)	16.3 (±4.1)	8.2 (±2.3)				
	11 Apr 2003	318 (±102)	26.7 (±5.1)	17.1 (±4.5)	9.2 (±2.9)				
	5 Mar 2003	297 (±100)	26.2 (±5.1)	16.6 (±4.5)	8.5 (±2.9)				
Chan Forecast (±SD)	24 Jun 2003	-	26 (±3)	15 (±2)	-				
Chan Polecast (LSD)	24 Apr 2003	-	26 (±3)	16 (±2)	-				

